Fifteen-minute consultation: An overview of major incidents

Stefani Widya, 1 Rebecca Hewitson, 2 Tulsi Patel, 3 Damian Roland, 3 Christopher Dadnam © 3

¹General Paediatrics, Leicester Royal Infirmary, Leicester, UK ²Paediatric Emergency Department, Cardiff and Vale University Healthboard, Cardiff,

³Paediatric Emergency Department, Leicester Royal Infirmary, Leicester, UK

Correspondence to

Dr Christopher Dadnam, Paediatric Emergency Department, Leicester Royal Infirmary, Leicester LE2 7DX, UK; chmdadnam@hotmail.com

Received 26 March 2022 Accepted 27 May 2022 **Published Online First** 15 June 2022

ABSTRACT

Major incidents are rare but require a large amount of preparation, co-ordination and communication across different emergency services and specialities. This ensures that casualties are efficiently managed within the constraints of limited clinical resources. This article aims to provide a brief understanding of what constitutes as a major incident, how it is declared and the chain of command in communication and action, focusing specifically on the paediatric process. It also aims to highlight important considerations that could potentially be missed (eg, the mental health impact, forensic evidence and so on).

can be classified as a major incident.

Major incidents can range from transportation accidents, natural disasters to acts of terrorism or violence. Approximately 14000 people from across the UK attended a concert at the Manchester Arena in 2017 when it was bombed killing 22 people (including children). Over 1000 were physically injured with many more suffering psychological and emotional trauma. Other forms of major incidents may have a smaller media presence but can be just as impactful. In 2020, the Centre for Research on the Epidemiology of Disasters recorded a total of 389 climate-related disasters resulting in 15080 deaths worldwide. These will likely become more common globally with an increase in the frequency of extreme weather events.3

It is a government regulation that all hospitals must have a mass casualty incident plan and provide training and learning exercises to its staff to be prepared for a major incident.4

AN OVERVIEW OF A MAJOR INCIDENT

As these events are uncommon, it is understandable to have a heightened level of concern and anxiety around what is expected from us and what happens throughout this process. Being preinformed is vital as the quality of our response can be critical to the outcomes for patients.

The specialist nature of paediatric care can mean that even a relatively small number of paediatric casualties can quickly overwhelm the usual capacity of a local health system. These are stressful, busy and emotionally challenging events, and being prepared may improve your ability to cope in the moment. Co-ordinating an effective response to a major incident requires everyone across the healthcare system to participate and this may mean that you could be asked to perform tasks outside of your routine practice. Attending major incident training and reviewing local guidelines will invaluably prepare you for such situations. Some hospitals will also have additional plans for managing mass paediatric casualty events.

This consultation will aim to further discuss how a major incident can be managed effectively through this specific paediatric event:

Case scenario

During your shift you hear from the nurse in-charge that a local school bus, with 20 children on-board, has collided with a truck on the motorway. Your hospital is the local unit and emergency services have arrived at the scene, with multiple causalities being reported.

INTRODUCTION

Major incidents are defined as specific events (or series of events) causing multiple casualties which exceeds the normal capacity of the NHS and/or other emergency services. A major incident tends to involve many people with varying degrees of severity of injuries and usually occurs in a sudden manner. However, numerous smaller-scale events in combination or anything which could overwhelm capacity

To cite: Widya S, Hewitson R, Patel T, et al. Arch Dis Child Educ Pract Ed 2023;108:236-241.

@ Author(s) (or their employer(s)) 2023. No

Check for updates

commercial re-use. See rights

and permissions. Published

by BMJ.




Figure 1 A major incident can be declared through an external source or internally by the hospital itself (usually by the ED consultant who raises it to the Senior Executive).⁶

WHO DECIDES IF THIS SITUATION IS A MAJOR INCIDENT?

Major incidents can be versatile in clinical presentations. in many different situations that can potentially overwhelm current services/capacity can be considered as a major incident. A major incident can be declared within or outside of a secondary care setting (figure 1).

When a major incident has been confirmed, the status shifts to 'Declared—Activate Plan'. This sets in motion a whole cascade of pre-agreed actions, outlined in the local major incident plan.

HOW IS KEY INFORMATION COMMUNICATED?

Given the scale of this event, critical information relayed between the local/regional hospitals, emergency services and the Regional/National Command Centres (set up for individual incidents) needs to be succinct and informed. The current structured format indicates time, location, type of incident, hazards, approximate number and severity of casualties, in which emergency personnel are already at the scene and if more or different expertise is needed. This is summarised in the acronym METHANE (see table 1). This information should be recorded in a place where staff can see and update it in the hospital (eg, white board in ED resus). As shown, our incident has also been relayed in this manner.

HOW IS THIS INCIDENT BEING COMMANDED, CONTROLLED AND CO-ORDINATED?

Our major incident will draw the attention of multiple emergency services including police, fire service, paramedics and motorway services. These teams individually provide excellent specialist

Table 1 A METHANE approach to the example scenario (adapted from 'Clinical Guidelines for Major Incidents and Mass Casualty Events' Version 2, September 2020, NHS)⁶

METHANE	Specific questions	Example: School bus crash
Major Incident	Has a major incident been declared? Record the date and time of 'standby' and 'declared' status.	3 March 2022 10:20 hours Major incident declared School bus vs Truck
Exact location	What is the exact location of the incident?	M1 Junction 11
Type of incident	Fire/Flood/Blast/Explosion/ Gunshots/Building collapse/ Chemical/Nuclear /Biological/Radiation	Road traffic accident
Hazards	What hazards or potential hazards can be identified?	Fuel—potential for fire/explosion Weather— hypothermia
Access	What are the best routes for access and exit to the hospital site?	ETA: 30 min Access routes— no further information
Number of casualties	How many casualties are there, and what condition are they in?	20 children (age 10–11) 6 adults Prehospital triage score: 1: 3 patients 2: 5 patients 3: Unknown 4: 3 patients Remaining: Uncategorised
Emergency services	Which, and how many, emergency services and personnel are required or are already on the scene?	On scene: 2 police cars 3 ambulances

Figure 2 Three-tiered command and control structure to a major incident (left) and JDM from JESIP (right). Adapted from JESIP.¹⁵ JDM, joint decision model; JESIP, Joint Emergency Services Interoperability Programme.

services; however, effective collaboration is critical to successfully manage a major incident. JESIP (Joint Emergency Services Interoperability Programme) was established in 2012 to ensure the emergency responders are trained and exercised to work together as effectively as possible, at all levels of command.⁵

During any major incident there is a three-tiered command structure; Gold (strategic), Silver (tactical) and Bronze (operational), for each emergency service responding. Gold and silver commanders are not typically located on scene, rather they formulate and strategise the wider implications of a major incident (allocation of resources and media information). Bronze (operational) commanders, directly and collaboratively work at the incident site with their respected staff members. These operational commanders will appoint a lead agency and communicate using the joint decision model throughout the incident to ensure a coordinated and informed response (figure 2).

For our incident, given the number of potential casualties, it is important that multiple hospital sites are activated, as attendees could quickly overwhelm a single unit. This overview of resource allocation will help ensure the injured children and adults, at the incident, are diverted to the most appropriate centres based on the extent of their injuries while continuing to manage the normal daily influx of admissions.

Most hospitals will have 'action cards' which are handed out to key members of staff to allocate

roles and duties. These cards normally state your allocated role at the top, followed by a checklist of responsibilities and actions (table 2, box 1).

Box 1 Example of tasks paediatricians might be involved in

As paediatricians, depending on our current role/ rotation, given the scale of potential influx, we might be involved in:

- Discharging current inpatients.
- Preparing the ward to receive patients.
- More direct assistance in the emergency department (eg, reviewing medical patients and/or helping paediatric and adult emergency clinicians manage the influx of minor and major injuries).

WHO IS TRIAGING THESE PATIENTS?

On scene, the ambulance service is responsible for triaging and escorting of all casualties to the hospital safely, if indicated. The triaging system they use consists of four categories (table 3).

National guidance suggests the use of Jump-START, a dedicated triage tool for children under 12 years (figure 3). However, a study conducted by Malick *et al* looking at retrospective data in a trauma registry suggested that JumpSTART had a low sensitivity in predicting P1 status. Instead, using the Battlefield Casualty Drills triage system universally for all patients could simplify processes, minimise confusion and allow for standardised training. Standardised training and application of a universal triage system could also simplify the process and minimise confusion. Figure 4 further

 Table 2
 Examples of 'action cards' that might be used in a major incident

major in	cident				
Action Card No 45—Me	d edical Co-ordinator in Paediatrics				
Job title	Paediatric Consultant or Registrar				
Base	Paediatric Hospital				
Report to	Paediatric Hospital Co-ordination Team Leader (Handset 1251)				
Actions to	be completed when incident STANDBY				
1	Liaise with Paediatric Surgical Co-ordinator to consider if any current or planned activities should be cancelled or postponed				
2	Consider retaining staff if a shift is about to end (until more information is available)				
Actions to be completed when incident DECLARED					
3	Go to Level 4 Meeting Room, Paediatric Hospital for status update and collect tabard and handset				
4	Put on the tabard 'Medic Co-ordinator'				
5	Collect handset 1253				
6	Go to Seahorse ward to undertake a paediatric discharge round—identify inpatients who can be treated elsewhere/discharged/transferred				
7	Escalate any difficulties or resource requirements to the Paediatric Co- ordination Team Leader on handset 1251				
8	Record all your actions and decisions with times on your decision log				
Actions to	be completed when incident STAND DOWN				
9	Return decision log, tabard and handset to level four meeting room				
10	Participate in hot debrief in level four meeting room				
Action Caro No 23—Pa	d ediatric Emergency Department Major (P2) Doctor				
Job title	Paediatric Consultant or Registrar				
Base	PED Majors (P2)				
Report to	Paediatric Emergency Department Co-ordination Team Leader (Handset 1342)				
Actions to	be completed when incident STANDBY				
1	Make a plan for all of your patients to leave the Emergency Department (admit or discharge)				
2	Continue seeing new patients as normal				
Actions to	be completed when incident DECLARED				
3	Go to Emergency Department Seminar Room for status update and collect tabard and handset				
4	Put on the tabard 'P2 Paediatrics'				
5	Collect handset 1350				
6	Assist with the departure of all patients from the Paediatric Emergency department—admit, discharge or send to alternative service				
7	Assess and treat patients in PED Majors				
8	Escalate any difficulties or resource requirements to the Paediatric Co- ordination Team Leader on handset 1342				
9	Keep incident notes folder updated for each patient with plan and likely destination				
10	Make sure every patient going to the ward has a clear plan and handover summary				
11	Make sure every patient being discharged has a GP letter mentioning that the patient was involved in the major incident				
12	Record all your actions and decisions with times on your decision log				
Actions to	be completed when incident STAND DOWN				
13	Return decision log, tabard and handset to Emergency Department Seminar Room				
14	Participate in hot debrief in Seminar Room				
15	Return to normal system of seeing patients in the majors area				

elaborates on NHS triaging system for major incidents and mass casualty events, highlighting differences in adult and paediatric triage.

WHO SHOULD ATTEND THE HOSPITAL TO HELP WITH A MAJOR INCIDENT?

Following this incident, your local major incident plan would have been executed, including specific

Table 3 Categorising children based on the triage criteria to their appropriate priority level. adapted from 'clinical guidelines for major incidents and mass casualty events' version 2, September 2020, NHS⁶

These are the most unwell or severely injured patients but for whom treatment and transfer can be lifesaving for example, major haemorrhage, open chest wounds	
Injured and unable to walk but can wait a short time for treatment for example, conscious with a head injury	
Minor fractures or lacerations	
Dead or obviously dying	

instructions for switchboard about who to contact. If you have learnt about the incident from colleagues or the media, it is important not to attend the hospital unless you have been asked to do so or would normally be on shift. Remember, the response to a major incident and the additional work in returning to normal afterwards can take days or weeks and it is important that staff are kept in reserve for this.

WHAT ELSE SHOULD BE CONSIDERED DURING A MAJOR INCIDENT?

Forensic evidence

It is important for the police to investigate how the incident unfolded. Any personal belongings or clothing removed from patients should therefore be minimally handled and while wearing gloves, before placing in individual plastic bags and keeping a note of what has been removed and where it is stored. As with safeguarding incidents, medical photographers may also be present to record injuries or body maps used to depict injuries.

Documentation

Documentation is crucial for recording what decisions were made and why, including the date, time and location of the patient. This is critical for individual patient care and because it may be used as evidence in a subsequent inquiry. All patients will be given temporary hospital numbers on arrival, and these must continue to be used until they are on the ward, even if their real identity is known later.

Mental health impact

As with any other major incident or traumatic life events, staff involved are likely to be impacted emotionally. Effective 'hot debriefs' immediately following the incident, provide a safe space to reflect on the incident: answering any concerns and highlighting any immediate lessons that have arisen. Poorly executed debriefs have the potential to cause more psychological harm. Subsequent formal multiagency debriefs following the event provide opportunities to review the JESIP principles applied, what worked well and what could be improved for future incidents. ¹⁰

There is an obvious immediate impact for children and families who have been involved in a major incident and responding to their psychological distress with compassion in the acute phase is a key part of the emergency response.¹¹ We also need to prepare families for the ongoing impact following discharge. Children involved in any major incident can be affected in several ways, and it is important to talk to families about what to expect (see box 2 for examples). Parents can support children by encouraging a return to their routine (including school) and being available to talk about the event and to listen to their child's feelings. An information leaflet for members of the public about dealing with the mental health impact of a major incident can be found in the NHS document 'Clinical Guidelines for Major Incidents and Mass Casualty Events'.

On arrival into hospital, a second triage is then performed. This can be emotionally challenging to triage patients to priority 4. The use of objective triage tools can help staff to make these difficult decisions. In these cases, it is important to remember that, although incredibly difficult, finite staff and resources should be prioritised for patients that are likely to survive given any intervention. Children who have been triaged will have a tag with their triage category. Patients for whom a trauma tourniquet has been applied to manage lifethreatening limb haemorrhage should have a 'T' and the time it was applied written on their forehead (and/ or over the tourniquet strapping) to minimise complications from prolonged application.

Box 2 Examples of how a child can be affected following a major incident

Some examples of what a child can present with due to the psychological distress following a major incident:

- Nightmares.
- Difficulty sleeping.
- Difficulty concentrating.
- ▶ Physical symptoms (eg, headaches or abdominal pain).
- Regression of previous skills (eq, toileting).

Media interest

There is likely to be media interest in the incident in which you have been involved and there may be members of the press attending the hospital to get information. Be wary of the press' attempts to obtain information or access the hospital and do not offer any interviews or comments of your own. Avoid posting anything on social media about the incident. This is crucial because of the potential impact on friends or family of those involved and also to ensure that any police investigations into the incident are not jeopardised.

Training and exercises

Who would have thought that *you* would be managing a major incident today? Uncertainty is the greatest




Figure 3 JumpSTART triage used to triage children at the scene of casualty. NHS. Clinical Guidelines for Major Incidents and Mass Casualty Events (Internet). 2nd ed. NHS England; 2020. Available from: https://www.england.nhs.uk/wp-content/uploads/2018/12/B0128-clinical-guidelines-for-use-in-a-majorincident-v2-2020.pdf (accessed 19 February 2022).⁶

challenge in the face of a mass casualty incident. It therefore becomes more difficult to plan and distribute critical resources. Although there will never be a point in time that there is sufficient information in a mass casualty incident, there are ways of mitigating the uncertainty including, preparation for such large-scale events through training. This can involve a variety of methods: discussion-based exercises such as workshops, seminars and lectures to operation-based exercises that simulate a scenario, drills and more recently the use of virtual reality. The seminary incident.

Making sure that you have read through your trusts local major incident plan will leave you feeling more prepared in the event of one taking place. They are lengthy documents and not intended to be read once a major incident has been declared!

Training exercises should aim to test different aspects of the response system including the clinical issues, command and leadership tactics and the

Emergency department triage (paediatric <12 years)

	erformed at the sce ioritise patients for	ene of the incident evacuation/transport	Secondary triage Performed to re-evaluate a patient after primary triage has been completed Typically done once the patient arrives in hospital Can also be done at an alternative care site, casualty clearing station or if time on scene is prolonged	
Priority 1	Immediate	Severely ill/injured but treatable. Able to be saved with relatively quick treatment and transport eg severe bleeding, sepsis, open chest or abdominal wounds, severe respiratory distress, emotionally uncontrollable.		
Priority 2	Delayed	Injured/ill and unable to walk on their own; potentially serious injuries/illnesses but stable enough to wait a short while for medical treatment eg burns with no respiratory distress, spinal injuries, moderate blood loss, conscious with a head injury.		
Priority 3	Minor	Minor injuries/illnesses that can wait for a longer period of time for treatment eg minor fractures, minor bleeding or minor lacerations.		
Priority 4	Expectant	Dead or obviously dying. May have signs of life but injuries are incompatible with survival eg cardiac arrest, respiratory arrest with a pulse*, massive head injury. It can be emotionally challenging to tag a child as expectant/deceased. Resist the tendency to assign a higher triage category to paediatric patients just because they are children. Sing an objective triage tool during a major incident can provide emotional support for staff forced to make these decisions for children.		
		* In children, typically respiratory failure precedes circulatory failure. If a child is apnoeic but has a pulse, a brief trial of ventilations, may 'jumpstart' their respirations (trial: five rescue breaths)		
Main differer	nces between adu	lt and paediatric triage		
 In children, if positioning the airway does not restart ventilation, then give a trial of ventilation, as this may restart spontaneous ventilation. In adults, there is no trial of ventilation and the adult casualty is tagged expectant or dead. 		ventilation, as this may on. In adults, there is no trial	The JumpSTART® paediatric triage MCI triage tool (usually shortened to JumpSTART®) is a variation of the simple triage an rapid treatment (START) riage system. Both systems are used to sort patients into categories at mass casualty incidents (MCIs).	
dead. In children, only peripheral pulses should be used to assess circulation. In children, APU is used to assess mental status, not ability to follow commands.		sess mental status,	However, JumpSTART® was designed specifically for triaging children in disaster settings. Though JumpSTART® was developed for use in children from infancy to age 8, where age is not immediately obvious, it is used in any patient who appea to be a child (patients who appear to be young adults are triage).	
Non-ambulatory children include:			using START or other adult triage systems).	

Figure 4 Further elaboration on triaging system from NHS. Clinical Guidelines for Major Incidents and Mass Casualty Events (Internet). 2nd ed. NHS England; 2020. Available from: https://www.england.nhs. uk/wpcontent/uploads/2018/12/B0128-clinical-guidelines-for-use-in-a-major-incident-v2-2020.pdf (accessed 19 February 2022).⁶

communication system. The aim of these exercises is to identify the strengths and limitations of the current system. As shown however, each incident is unique, and every plan will inevitably have to be adapted to accommodate the situation.

CONCLUSION

Major incidents are rare; however, they require a great deal of coordination, adaptability and communication between multiple service providers. This multiagency collaboration ensures that all patients are treated appropriately, while accounting for the limited resources and situational uncertainty. Frequent effective training (involving all the relevant departments) is an essential component that is required for each hospital, to ensure their staff are familiar with their roles in a major incident. Moreover, particular attention and training, through JESIP and major incident resources, towards paediatric major incidents will help to build a network of systems that is more adaptable and ready to manage such events.

Correction notice This article has been updated since it was first published. Table 2 and boxes 1 and 2 have been moved closer to their citations.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Commissioned; externally peer reviewed.

RCID iD

Christopher Dadnam http://orcid.org/0000-0001-8626-4194

REFERENCES

- 1 JESIP. The Kerslake report: an independent review into the preparedness for, and emergency response to, the Manchester arena attack on 22ndMay 2017, 2017. Available: https://www.jesip.org.uk/uploads/media/Documents%20Products/Kerslake_Report_Manchester_Are.pdf [Accessed 13 Feb 2022].
- 2 Centre for research on the epidemiology of disasters; un office for disaster risk reduction. 2020: the non-COVID year in disasters — global trends and perspectives. Brussels. un 2021. Available: https://www.undrr.org/publication/2020-non-covid-year-disasters [Accessed 13 Feb 2022].
- 3 Intergovernmental panel on climate change; climate change 2022 impacts, adaptation and vulnerability. IPCC, 2022. Available: https://report.ipcc.ch/ar6wg2/pdf/IPCC_AR6_WGII_FinalDraft_FullReport.pdf [Accessed 13 Mar 2022].
- 4 Moss R, Gaarder C. Exercising for mass casualty preparedness. *Br J Anaesth* 2022;128:e67–70.
- 5 Critical incident management. Home Office [Internet], 2021. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1013914/national-critical-incident-management-guidance-v13.0-ext.pdf [Accessed 13 Feb 2022].
- 6 NHS. Clinical Guidelines for Major Incidents and Mass Casualty Events [Internet]. 2nd ed. Available: https://www.england.nhs.uk/wp-content/uploads/2018/12/B0128-clinical-guidelines-for-use-in-a-major-incident-v2-2020.pdf [Accessed 19 Feb 2022].
- 7 Malik NS, Chernbumroong S, Xu Y, et al. The BCD triage sieve outperforms all existing major incident triage tools: comparative analysis using the UK national trauma registry population. EClinical Medicine 2021;36:100888.
- 8 Malik NS, Chernbumroong S, Xu Y, et al. Paediatric major incident triage: UK military tool offers best performance in predicting the need for time-critical major surgical and resuscitative intervention. EClinicalMedicine 2021;40:101100.
- 9 Cooper S, Winton M, Farrington-Exley J. Fifteen-minute consultation: time out as an alternative to toxic debrief. *Arch Dis Child Educ Pract Ed* 2020;105:270–5.
- Wolfe HA, Wenger J, Sutton R, et al. Cold Debriefings after in-hospital cardiac arrest in an international pediatric resuscitation quality improvement collaborative. Pediatr Qual Saf 2020;5:e319.
- 11 Desmond M, Schwengel D, Chilson K, *et al.* Paediatric patients in mass casualty incidents: a comprehensive review and call to action. *Br J Anaesth* 2022;128:e109–19.
- 12 Tallach R, Brohi K. Embracing uncertainty in mass casualty incidents. *Br J Anaesth* 2022;128:e79–82.
- 13 Lefevre-Scelles A, Barege V, Roussel M, *et al.* Development and evaluation of a mass casualty medicine simulation method using figurines. *Br J Anaesth* 2022;128:e210–2.
- 14 Skryabina EA, Betts N, Reedy G, et al. The role of emergency preparedness exercises in the response to a mass casualty terrorist incident: a mixed methods study. Int J Disaster Risk Reduct 2020;46:101503.
- 15 JESIP Resource Manual [Internet]. JESIP, 2022. Available: https://www.jesip.org.uk/resource-manual